arXiv:1211.4082v1 [cond-mat.stat-mech] 17 Nov 2012

Keywords: phase transition,nonlinear model, coupling

Model of phase transitions with coupling of the order parameter and its gradient

B. I. Lev and A. G. Zagorodny
Bogolyubov Institute for Theoretical Physics, NAS Ukraine, Metrolohichna 14-b,
Kyiv 03680, Ukraine. E-mail blev@bitp.kiev.ua, Tel/Fax-044-521-31-63

(Dated: November 20, 2012)

A model of phase transitions with coupling between the order parameter and its gradient is proposed. This nonlinear model is shown to be suitable for the description of phase transitions accompanied by the formation of spatially inhomogeneous distributions of the order parameter. Exact solutions of the proposed model are obtained for the special cases which can be related to the spinodal decomposition or cosmological scenario. Such solutions manifest crucial dependence on the coupling constant.

PACS numbers: 64.60.Cn, 75.40.Mg

Within the context of the general theory of phase transitions, a system treated as a continuous medium is assumed to have a ground state which can always be described in terms of the order parameter. Usually, the ground state is described by a definite value of the order parameter that is independent of the point of the elastic continuum. The order parameter associated with the phase transitions can have various geometrical presentations, for example, a scalar field in the case of condensed matter [1], or a fundamental scalar field in the field theory [2]. To introduce the order parameter that determines a stable state of the condensed matter, we have to consider probable deformations of the field distribution, in particular, the disordered configuration of the ground state. Such distributions of the order parameter can be observed experimentally, this issue requires an appropriate theoretical description. Today there exist many phenomenological models which are declared as those that can give fairly general description of phase transitions and the behavior of spatial distributions of order parameters both before and after the phase transition. To these well-known phenomenological models one can attribute the Landau theory of phase transitions [1] and the gradient theory of phase transitions [3],[4]. In the case of the first model, the free energy of the system under consideration is described in terms of the order parameter while in the second case the description is given in terms of the order parameter gradient. Both models provide an efficient description of many details of phase transitions, in particular, the post-transition behavior of spatial distributions of the order parameter.

The purpose of the present paper is to show that the standard model of phase transitions can be unified with the gradient theory by introducing in the standard model the coupling between the order parameter and its gradient. In this case we get rid of the requirement to describe only phase transitions with homogeneous distributions of the order parameter. Namely, problems of this kind arise in modern physics, including the spinodal decomposition [5] and the alternative cosmological model that takes into account the formation of inhomogeneous distributions in the fundamental scalar field [2]. As it was shown in [5], the coefficients before the quadratic terms of the order parameter and its gradient can change their signs at the instability thresholds and thus some decomposition scenario becomes possible. This can result in the transition from the disordered state to modulated, patterned, or ordered-patterned states. In particular, in the case of cosmological scenario, it would be highly desirable to describe the generation of spatially inhomogeneous states associated with the specific behavior of the coefficients in terms of the expansion of the model potentials. When the system described by a scalar spatially-dependent order parameter is rapidly quenching from a homogeneous high-temperature phase to the phase-coexistence region, we have the spinodal decomposition. The decrease of the temperature can make the reason for the formation of a new bubble phase in the cosmological model. Usually, the theoretical analysis of the spinodal decomposition and spatial formation of the fundamental scalar field is based on the standard model. In this case, however, the presentation for the free energy contains no terms that would take into account the interaction between the order parameter and its gradient though this interaction can lead to the restriction of the governing parameter.

Any deformation of the order parameter requires additional energy and thus imposes restrictions on probable deformations. In other words, not all deformations can be realized in system. In the general case the requirement of consistency of the known solutions of the standard model with the generalized description can be provided by an appropriate choice of the coupling between the order parameter and its gradient. As it was mentioned above, the phase transitions associated with the system with broken continuous symmetry can be described in terms of the relevant order parameter. In particular, according to the Landau theory, the free energy density can be presented in terms of the order parameter as given by

$$F = C\left(\nabla\varphi(\mathbf{r})\right)^2 + W(\varphi(\mathbf{r})) \qquad (1)$$

where $\varphi(\mathbf{r})$ is the order parameter, $W(\varphi(\mathbf{r}))$ describes the order-parameter dependence of the free energy that is assumed to be known, and C is a constant.

In the well-known standard model of phase transitions we have

$$W(\varphi(\mathbf{r})) = -\frac{1}{2}A\varphi^2(\mathbf{r}) + \frac{1}{4}B\varphi^4(\mathbf{r}). \tag{2}$$

With the dimensionless variable $\varphi^2(\mathbf{r}) = \frac{B\varphi^2(\mathbf{r})}{A}$ being introduced, the standard dimensionless free energy reduces to the form given by

$$f = l^2\left(\nabla\varphi(\mathbf{r})\right)^2 + ((\varphi(\mathbf{r}))^2 - 1)^2 \tag{3}$$

or

$$f = l^2\left(\nabla\varphi(\mathbf{r})\right)^2 + ((\varphi(\mathbf{r}))^2 - 1)\varphi^2(\mathbf{r}) \tag{4}$$

with the potential being written in the standard form and $l^2 = \frac{C}{A}$ being the characteristic length. It should be noted that in this presentation the dimensionless free energy depends on the relation between A and B and its sign. Making use of this expression for the free energy density in mean field approximation, we can find the spatial distribution of the order parameter and thus to describe the properties of the new states which can be formed after the phase transition. The first presentation of the free energy density generates the Euler-Lagrange equation for the stationary distribution of the order parameter that is given by

$$l^2(\nabla)^2\varphi(\mathbf{r}) = 2\varphi(\mathbf{r})(\varphi^2(\mathbf{r}) - 1) \tag{5}$$

and has a topological soliton solution $\varphi = tanh(\frac{r}{l})$. In the second case we can obtain another form of the Euler-Lagrange equation, i.e.,

$$l^2(\nabla)^2\varphi(\mathbf{r}) = \varphi(\mathbf{r})(1 - 2\varphi^2(\mathbf{r})) \tag{6}$$

. The latter equation has a different solution, $\varphi = sech(\frac{r}{l})$, which contains no topological singularities. This example shows that different coefficients in the expansion of the free energy result in different spatial distributions of the order parameter which can be observed experimentally.

In the case of a system with the gradient of the order parameter, it is likely to be reasonable to introduce, in the free-energy functional, a term responsible for the probable interaction between the order parameter and its gradient. This coupling can regularize possible perturbations of the order parameter and thus confine the spatially inhomogeneous state of the system. This means that not any possible deformation of the order parameter can exist in the deformed matter. Any defect introduces additional energy and not any deformation can contribute to the spatial distribution of the order parameter.

The main idea of the present contribution is to generalize the phenomenological theory of phase transitions by introducing the interaction between the order parameter and its gradient. Such generalization is reasonable in view of both mathematical and physical arguments. With regard to the above assumptions, the free energy can be presented in the form

$$\begin{gathered} f = a\left(|D\varphi(\mathbf{r})|\right)^2 + b((|\varphi(\mathbf{r})|)^2 - 1)^2 + \\ + c\varphi^2(\mathbf{r})\left(D\varphi(\mathbf{r})\right)^2 \end{gathered} \tag{7}$$

where operator $D = l\nabla$ is introduced, with the length of change of the order parameter; l, a,b, and c are parameters describing the influence of the gradient of the order parameter, the order parameter, and the coupling between the order parameter and the order parameter gradient. If the coefficient c is equal to zero, then we return to the standard theory of phase transitions or the theory of fundamental scalar field. Nevertheless, these considerations demonstrate the need to treat the variation problems for the functional in various representations. We prove that, as the relaxation parameter vanishes, the families of such fields with finite free energy are compact and have many solutions which describe spatially inhomogeneous distributions of the order parameter and possible topological structures of the new phase. Our proof is based on the analysis of the extremum of the free energy density. The method also yields an Euler-Lagrange equation for the limit of the defect structure measure. This measure also satisfies the cancellation properties depending on its local regularity, which seems to indicate several levels of singularities in the limit. We show that the same solution can be obtained in other interpretation. To do this, we have to analyze the behavior of the free energy density. The free energy minimum satisfies the Euler-Lagrange equation

$$\begin{gathered} (a + c\varphi^2(\mathbf{r}))D^2\varphi(\mathbf{r}) + c\varphi(\mathbf{r})(D\varphi(\mathbf{r}))^2 \\ +2b\varphi(\mathbf{r})(1 - \varphi^2(\mathbf{r})) = 0 \end{gathered} \tag{8}$$

Let us consider several possible solutions of the Euler-Lagrange equation for various combinations of coefficients in the one-dimensional case, when $D\varphi = \frac{d\varphi(\mathbf{r})}{dx} = \dot{\varphi}$. The Euler-Lagrange equation can be rewritten in a simple form, i.e.,

$$\ddot{\varphi} + g(\varphi)(\dot{\varphi})^2 + h(\varphi) = 0 \tag{9}$$

where $g(\varphi) = \frac{c\varphi}{a + c\varphi^2}$ and $h(\varphi) = \frac{2b\varphi(1-\varphi^2)}{a + c\varphi^2}$ If a new variable $u(\varphi) = (\dot{\varphi})^2$is introduced, then the equation thus obtained can be reduced to the Bernoulli equation, i.e.,

$$u(\acute{\varphi}) + 2g(\varphi)u(\varphi) + h(\varphi) = 0 \tag{10}$$

where $u(\acute{\varphi})$ implies derivation with respect to φ. The general solution of the Bernoulli equation in our case has the form

$$u(\varphi) = \frac{1}{a + c\varphi^2}(s + b(1 - \varphi^2)^2) \tag{11}$$

where s is the integration constant which in the general case can be put equal to zero. From this solution of the

Bernoulli equation we obtain that the spatial behavior of the order parameter is described by the relation

$$\frac{d\varphi}{dx} = \frac{b^{\frac{1}{2}}(1-\varphi^2)}{(a+c\varphi^2)^{\frac{1}{2}}} \tag{12}$$

which yields an expression for the spatial distribution of the order parameter given by

$$\begin{aligned} x = (\tfrac{c}{b})^{\frac{1}{2}} Arsh((\tfrac{c}{a})^{\frac{1}{2}}\varphi) + (\tfrac{a+c}{b})^{\frac{1}{2}} Arsh((\tfrac{a}{c})^{\frac{1}{2}} \\ + \tfrac{1}{2}(\tfrac{a+c}{b})^{\frac{1}{2}} Arsh((\tfrac{2\varphi\sqrt{(a+c)(a+c\varphi^2)}}{a(\varphi^2-1)} \end{aligned} \tag{13}$$

The behavior of solutions for different signs of the coupling constant is shown in Fig.1. Thus, we can obtain from this solution all the well-known standard solutions. For $c = 0$, we find that



FIG. 1: Different solutions for various values of the coupling constant

$$x = \frac{1}{2}(\frac{b}{a})^{\frac{1}{2}} Arth(\frac{2\varphi}{\varphi^2-1}) \tag{14}$$

or $\varphi = th(\frac{a}{b})^{\frac{1}{2}}x$ that fully recovers the solution of the standard theory of phase transitions, or the behavior of the fundamental scalar field in the standard cosmological model. If the coupling constant becomes negative, then the solution crucially changes. In the case when the $c = -a$, we obtain a periodic solution $\varphi = sin(\frac{b}{a})^{\frac{1}{2}}x$. This solution is not trivial in the physical sense. Within the context of this solution we can propose a new scenario of phase transitions. Namely, we can predict the existence of various spatial distributions of the order parameter after the formation of the new phase. In order to make such a prediction we have to know (or postulate) the temperature dependence of the coupling constant. If the coupling between the order parameter and its gradient is included into the free energy with positive value, we can observe an inhomogeneous distribution of the order parameter or the fundamental scalar field without topological singularities. If the coupling constant changes as temperature changes and if its temperature-dependence is critical, $c = c(T - T_s)$, where T_s is the new critical value of temperature for the coupling constant equal to zero, then we can observe solution of the standard model of phase transitions, $\varphi = th(\frac{a}{b})^{\frac{1}{2}}x$ with the topological singularity. The further decrease of temperature can lead to the negative value of the coupling constant with the dependence $c = c(T - T_s) = -a$ and the topological solution transforms to the periodic solution $\varphi = sin(\frac{b}{a})^{\frac{1}{2}}x$, i.e., a regular solution without any singularities.

Assuming that the interaction between the order parameter and its gradient induces selection of possible deformations that depends on the coupling constant, we can obtain various spatial distributions of the order parameter. Such phase transitions can be observed in the experiments with decreasing temperature. This effect can be expected in the standard cosmological model, when we have evolution of the bubble formation (as a topological formation) to a new periodic phase of the fundamental scalar field without singularities. This effect should be observed in the experimental studies of the spinodal decomposition.

To conclude, we note that the model of phase transitions with coupling between the order parameter and its gradient can describe a new scenario of the spatial distribution of the order parameter after the critical point of the phase transition. In this model we obtain an exact analytical solutions and predict the experimental observation of the new behavior of the order parameter in the case of the spinodal decomposition or in the cosmological scenario in the field theory. It is shown that this nonlinear model can be used for the description of phase transitions accompanied by the formation of spatially inhomogeneous distributions of the order parameter for various systems. Such solutions crucially depend on the coupling constant. In the case of its negative value we can expect that the distribution of the order parameter should be periodic, but in the case of a positive coupling constant, we obtain solutions with topological singularities.

---

[1] L. D. Landau and E. M. Lifshiz *Field theory* Nauka, (1973).

[2] A. D. Linde, *Elementary particle physics and inflationary cosmology*, Horwood Academic, Switzerland,(1990).

[3] P. Aviles and Y. Giga, Proc.Center Math Analysis Australian National University,**12**,1 (1987),

[4] P. Aviles and Y. Giga, Proc. Royal. Soc. Edinburgh, **A129**, 1, (1999).

[5] S.B. Goryachev, Phys. Rev. Lett., **72**, 1815, (1994)

[6] B. S. DeWitt *Phys.Reports* **19C**, 295, (1975)

[7] P. Chaikin and T. C. Lubensky, *Principle of condensed matter physics*, Cambridge,(1998).
[8] P. G. de Gennes and J. Prost. The Physics of Liquid Crystals (2ed.,Clarendon Press, Oxford,1993).